

**Species composition and seasonal variation of phytoplankton  
in Himreen reservoir, middle of Iraq**

**Abbas Murtadeh Ismail**

**Diyala University, College of Education, Biology Department**

**Abstract:**

The Phytoplanktons of Himreen reservoir were studied during the period Feb. 1996 to Jan. 1997. A total of 98 algal species were identified, dominated by diatoms (69 species), followed by green algae (15 species), blue green algae (10 species) and the other taxa (4 species).

Diatoms were the most abundant group, followed by green and blue green algae and the dominant species were *Navicula cryptocephala*, *Nitzschia palee*, *Cymbella affinis*, and *Fragillaria ulna*. The number of species revealed clear differences and irregular seasonal variations at different stations of the study area.

## **Introduction:**

There are more than 66 water regulation projects in Iraq include dams, impoundments, reservoirs, flood regulation barrages (Saddalla, 1998)<sup>(1)</sup>. Himreen reservoir is one of the largest lentic aquatic system in Iraq. Only two ecological investigations have been published on this reservoir<sup>(1,2)</sup>. The reservoir received its water mainly from Diyala river which is the main and the earliest tributaries of river Tigris. It is considered one of major irrigation projects in Iraq. The present investigation was attempted to give further information on the phytoplanktons of Himreen reservoir qualitatively and quantitatively, as well as their seasonal variations.

## **Study Area:**

Himreen reservoir started operation since 1982, it is located about 40 km north east part of Diyala governorate, with about 374km<sup>2</sup> storage area with maximum depth of 35m, at about 53m above sea level<sup>(3)</sup>. The reservoir is subjected of sever water level fluctuation leading a short and long term effect on the available organisms. Two stations were selected for monthly sampling within the reservoir (Fig. 1).

## **Materials and Methods:**

Monthly surface water samples were collected during the period from February 1999 to January 2000 from two selected stations in Himreen reservoir (Fig. 1). At each station a Hydro-Bios plankton net of 20 $\mu$ m mesh size which was towed just beneath water surface about 15 minutes for qualitative study. Sedimentation technique was followed for quantitative study<sup>(4)</sup>, as cited by Kassim, et. al. (1999)<sup>(5)</sup>.

Diatoms were cleaned by using hot nitric acid and counted by microtransect method<sup>(6)</sup>, whereas the non-diatoms algae were counted by haemocytometer chamber and the species identification was followed references<sup>(7-10)</sup>.

### Results and discussion:

A total of 98 algal taxa was identified in Himreen reservoir during the study period. Out of the total species the diatoms was dominated (71.42%) followed by green algae (15.3%), blue green algae (10.2%), and the others (4.08%), results reveals that 50 sps. Are common among the two stations (table 1). The number of chlorophycean species exceed that of cyanophycean (table 2), this gives an indication that the water is clean which was reported earlier during examination of water characters<sup>(1)</sup>. The species richness in the studied area is less than that in Qadissia lake<sup>(5)</sup> and Habbaniya lake<sup>(11)</sup>. Diatoms was the most dominant group in lakes all Iraqi, such as Dokan lake<sup>(12)</sup>, Razzazah lake<sup>(13)</sup>, Qadissia lake<sup>(5)</sup> as well as marshes area<sup>(14)</sup> and Al-Saadi et. al.<sup>(15)</sup>. Only 4 sps. of Euglenophyceae as well as pyrrophyceae were identified in the studied area. Some of the identified species in the present study are originated mainly from benthic forms.

Similarly several species of phytoplankton in Iraqi lakes illustrated that, such as Razzazah lake<sup>(16)</sup>. Few genera in the studied area were represented by several species such as *Nitzschia* (16 sps.), *Cymbella* (6 sps.) and *Navicula* (8 sps.) table (2). Similarly these genera were also found in several species in Qadissia lake<sup>(5)</sup> and Habbaniya lake<sup>(11)</sup>, in the mean time few species appeared in more than 8 months such as *Cymbella affinis*, *Nitzschia palae*, *Fragillaria ulna*, and *Navicula cryptocephala*. The total cell number of phytoplankton was  $250.9 \times 10^3$  and  $300.4 \times 10^3$  cells/l in sts.1 and 2 respectively and dominated by diatoms also in both stations. On the other hand, the cell number was more in st.2 also along the studied period (Fig. 2). The dominancy of diatoms in the cell count was recorded in the studied lakes and marshes which indicated above. Results reveals that the algal bloom was observed in the Spring and Autumn seasons (Fig. 3). Spring is shown to be the season of the higher growth and diversity, this result is similar to other Iraqi inland waters<sup>(5,11,12)</sup>. As indicated above the species composition and density were higher at st.2 which may be explained due to the variation in some related environment factors which need more future research in the investigated area.

## References:

1. Saadalla, H. A. A. (1998). Ecological study on the effect of Himreen impoundment on the benthic and planktonic invertebrates of river Diyala, Ph. D. thesis (In arabic), Univ. Baghdad, 254 pp.
2. Sulaiman, N. I; Saadalla, H. A. A; and Ismail, A. M. (2001). Regulation influence of Himreen reservoir on phytoplankton in river Diyala, Iraq. *J. Environ. Stud.* 58:749-760.
3. Al-Saadi, H. A. M. (1986). Irrigation and drainage projects in Diyala province. Geographic study (In arabic), M. Sc. Thesis, Univ. Baghdad, 250 pp.
4. Martinez, M. R.; Chakroff, R. P. and Pantastico, J. B. (1975). Note on direct phytoplankton counting technique using the haemocytometer. *Phil-Agric*, 59:1-12.
5. Kassim, T. L.; H. A. Al-Saadi, A. A.; Al-Lami and Y. A. Alwan. (1999). Spatial and seasonal variations of phytoplankton in Qadisia lake, Iraq. *IACE Sci. J.* 1:99-111.
6. Hadi, R. A. M. (1981). Algal studies of the river USK, Ph. D. Thesis, Univ. Coll., Cardiff., U.K. 364 pp.
7. Desikachary, T. V. (1959). *Cyanophyta*. Acad. Press, London.
8. Germain, H. (1981). *Flore des Diatom' ees. diatomophyceae aedouces et samtres du Massif Armoricion et des contr, ees voisines d' Europe occidental-* Paris, Soc. Nouv. Ed. Boub'e 2, 444pp.
9. Hustedt, F. (1985). The pinnate diatoms: Strauss and Gramer on English translation of "Die kieselalgen, teil 2" with supplement by N. Jensen.
10. Prescott, G. W. (1973). *Algal of the Western Great Lakes area*. Dubuque, W. C. 13 rown. co, 977 pp.
11. Kassim, T. I; Al-Saadi, H. A; Salman, S. K. and Farhan, R. K. (2001). Species composition and seasonal variation of phytoplankton in Habbaniya Lake, Iraq. *Iraqi J. of Biology.* 1(1) 23-34.
12. Shaban, A. A. (1980). An ecological study on the phytoplankton of Dokan Lake. M. Sc. Thesis. Univ. Sulaimaniyah, Iraq.
13. Anon. (1983b). State and prospection of Fisheries in Razzazah lake. Final report. Poleservice cons. Engine. Warsaw-Poland. State Fisheries organization, Baghdad, Iraq.
14. Al-Mousawi, A. H.; H. A. Al-Saadi and F. M. Hassan. (1994). Spatial and seasonal variations of phytoplankton populations and

- related environment in Al-Hammar marsh, Iraq. Bas. J. Sci. 12(1):9-20.
15. Al-Saadi, H. A. and A. A. Al-Lami (1992). Seasonal variation of phytoplankton in some marsh areas in southern Iraq. S. Coll. Educ. For Women, Univ. Baghdad, 3:56-61. (In arabic).
  16. Al-Ghafily, A. K. (1992). Ecological study on phytoplankton in Razzazah lake, Iraq. M. Sc. Thesis. Univ. Baghdad, Iraq. (In arabic).




Table 1: List of algal taxa, total count (%) and appearance refers to number of months that the species was identified in studied stations.

Taxa	st.1	st.2	% cell Nu.	App.
<b>CYANOPHYCEAE</b>				
<i>Anabaena</i> sp.	+	+	0.79	3
<i>Chroococcus minor</i> (Kuetz.) Naegel:	+	+	0.69	3
<i>Chroococcus</i> sp.	-	+	0.20	2

<i>Merismopedia elegans</i> A. Br.	+	+	0.35	4
<i>Merismopedia glauca</i> (Ehr.) Naeg.	-	+	0.53	1
<i>Microcystis aeruginosa</i> Kuetz.	+	+	0.34	2
<i>Microcystis</i> sp.	+	+	0.16	1
<i>Oscillatoria tenuis</i> C.A.Agardh	+	+	0.81	5
<i>Oscillatoria</i> sp.	+	-	0.12	2
<i>Spirulina</i> sp.	+	+	0.3	2
<b>EUGLENOPHYCEAE</b>				
<i>Euglena</i> sp.	+	+	0.22	3
<i>Phacus</i> sp.	+	-	0.09	1
<b>PYRRHOPHYCEAE</b>				
<i>Peridinium cinctum</i> (Mell) Ehr.	+	+	0.11	2
<i>Peridinium</i> sp.	-	+	0.05	2
<b>CHLOROPHYCEAE</b>				
<i>Ankistrodesmus</i> sp.	+	+	0.44	3
<i>Chlamydomonas</i> sp.	+	+	0.51	4
<i>Chlorella vulgaris</i> Beyerinek	-	+	0.12	2
<i>Closterium</i> sp.	-	+	0.19	3
<i>Coelastrum reticulatum</i> (Dang.) Sen.	+	+	0.24	4
<i>Coelastrum</i> sp.	-	+	0.10	2
<i>Cosmarium</i> sp.	+	+	0.07	1
<i>Oedogonium</i> sp.	+	+	0.66	3
<i>Pediastrum duplex</i> Lager.	+	+	0.77	6
<i>Pediastrum simplex</i> (Meyen) Lemm.	+	-	0.41	3
<i>Pediastrum</i> sp.	+	+	0.11	1
<i>Scenedesmus bijuga</i>	+	+	0.55	4
<i>Spirogyra</i> sp.	+	+	0.33	3
<i>Tetradraeron minimum</i>	+	-	0.21	2
<i>Tetradraeron</i> sp.	+	+	0.56	3
<b>BACILLARIOPHYCEAE</b>				
"centrals"				
<i>Aulacosiera granulate</i> (Ehr.) Simo.	+	+	2.55	6
<i>Coscinodiscus lacustris</i> Grun.	-	+	0.33	5

<i>Cyclotella kuetzingiana</i> Thw.	-	+	0.41	4
<i>Cyclotella meneghiniana</i> Kuetz.	+	+	1.22	6
<i>Cyclotella ocellata</i> Panto.	+	+	3.01	6
<i>Cyclotella stelligera</i> Cl. El.	-	+	6.11	5
<i>Cyclotella</i> sp.	+	-	1.15	2
"pennales"				
<i>Achnanthes minutissima</i> Kuetz.	+	+	9.32	7
<i>Amphiprora alata</i> Kuetz.	-	+	3.7	3
<i>Amphora ovalis</i> Kuetz.	-	+	1.66	2
<i>Amphora veneta</i> Kuetz.	+	-	0.44	1
<i>Amphora</i> sp.	+	-	1.00	2
<i>Anomoeneis exilis</i> (Kuetz.) Cl.	+	+	3.14	4
<i>Bacillaria paxillifer</i> (Mull.) Hend.	+	+	3.3	4
<i>Caloneis</i> sp.	-	+	0.09	1
<i>Cymatoplura solea</i> de Br.	+	+	0.35	2
<i>Cymbella affinis</i> Kuetz.	+	+	10.65	8
<i>Cymbella amphicephala</i> Grun.	+	+	2.19	3
<i>Cymbella microcephala</i> Grun.	+	-	1.36	3
<i>Cymbella pusilla</i> Grun.	-	+	1.05	2
<i>Cymbella reutricosa</i> Kuetz.	-	+	0.17	1
<i>Cymbella</i> sp.	+	+	1.77	4
<i>Diatoma elongatum</i> (Lyng.) Agard	+	+	0.40	2
<i>Diatoma vulgare</i> Bory.	+	+	0.26	4
<i>Diploneis ovalis</i> Hislc.	+	-	0.41	3
<i>Diploneis</i> sp.	-	+	0.11	2
<i>Fragillaria acus</i> Kuetz.	+	-	0.24	2
<i>Fragillaria ulna</i> (Nitz.) Ehr.	+	+	6.66	9
<i>Fragillaria</i> sp.	+	-	0.66	2
<i>Gomphonema angustatum</i> (Kuetz.) Rab.	+	+	4.1	5
<i>Gomphonema</i> sp.	-	+	0.09	2
<i>Gyrosigma acuminatum</i> (Kuetz.) Rab.	+	+	0.56	5
<i>Gyrosigma spencerii</i> (W. Smith) Cl.	-	+	0.38	4
<i>Gyrosigma</i> sp.	+	+	0.28	3
<i>Hantzschia amphioxus</i> (Ehr.) Grun.	+	+	3.3	5
<i>Mastigloia smithii</i> Thw.	-	+	0.46	3
<i>Mastigloia</i> sp.	+	+	0.3	2



<i>Navicula anglica</i> Ralf.	+	-	0.29	2
<i>Navicula cryptocephala</i>	+	+	6.8	8
<i>Navicula cuspidata</i> Kuetz.	+	-	1.12	4
<i>Navicula gracilis</i> Ehr.	-	+	1.3	2
<i>Navicula pusilla</i> W. Smith	-	+	0.93	2
<i>Navicula radiosa</i> Kuetz.	+	-	0.30	4
<i>Navicula tuscula</i> (Ehr.) Grun.	+	+	1.11	5
<i>Navicula</i> sp.	+	-	2.3	6
<i>Nitzschia acicularis</i> (Ehr.) W. Smith	+	+	0.55	5
<i>Nitzschia amphibia</i> Grun.	-	+	0.86	4
<i>Nitzschia angustata</i> (W. Sm.) Grun.	+	+	0.66	4
<i>Nitzschia apiculata</i> (Greg) Grun	-	+	0.53	3
<i>Nitzschia dissipata</i> (Kuetz.) Grun	+	+	0.23	4
<i>Nitzschia fasciculata</i> Grun.	+	+	0.73	6
<i>Nitzschia frustulum</i> Kuetz.	+	+	0.81	5
<i>Nitzschia longissima</i> (Breb.) Ralf.	-	+	1.8	4
<i>Nitzschia lorenziana</i> Grun.	+	+	0.31	3
<i>Nitzschia microcephala</i> Grun.	-	+	0.16	2
<i>Nitzschia obtusa</i> W. Smith	+	-	0.21	3
<i>Nitzschia Palae</i> (Kuetz.) W. Smith	+	+	11.8	11
<i>Nitzschia sigma</i> (Kuetz.) W. Smith	-	+	0.12	2
<i>Nitzschia sigmoidae</i> (Ehr.) W. Smith	-	+	0.09	2
<i>Nitzschia tryblionella</i> Hantz.	+	-	0.36	1
<i>Nitzschia</i> sp.	+	+	2.1	3
<i>Pinnularia</i> sp.	-	+	0.07	1
<i>Rhoicosphaenia curvata</i> (Kuetz.) Grun.	+	+	0.81	4
<i>Stauronesis</i> sp.	+	-	0.02	1
<i>Surirella angustata</i> Kuetz.	+	+	0.24	2
<i>Surirella ovalis</i> de Breb.	+	+	1.33	3
<i>Surirella ovata</i> Kuetz.	-	+	0.56	2
<i>Surirella</i> sp.	-	+	0.21	3
<i>Tabellaria</i> sp.	+	+	0.07	1

Table 2: Number of identified species and genera of different algal classes in the studied stations

G.= genera      Sp.=species

Algal groups	St.1		St.2	
	G.	Sp.	G.	Sp.
Cyanophyceae	6	8	6	9
Euglenophyceae	2	2	1	1
Pyrrhophyceae	1	1	2	2
Chlorophyceae	7	13	10	13
Bacillariophyceae	21	46	24	54
Total Number	37	70	43	79

